

## Factors Affecting the Response to Erythropoietin in Hemodialysis Patients

العوامل المؤثرة على الاستجابة للريثروبويتين عند المرضى اللذين يخضعون لغسيل الكلى

By

Ali Abdulhasan Kamil

Supervisor

Dr. Hekmat Al-Akash

August 2020

### ABSTRACT

**Background:** Chronic kidney disease is associated with high risk of mortality and morbidity due to a multiplicity of factors. Erythropoietin stimulating agents are mainly used to correct anemia in patients with chronic kidney disease undergoing Hemodialysis.

**Objectives:** This study aimed to investigate the association of several factors (Kt/v, serum phosphorus, CRP, serum albumin, ESAs dose per week, ESAs type, iron level, number of HD session/ week, vitamin B12 level) with the erythropoietin response in patients with Hemodialysis.

**Methods:** A retrospective design was used to collect data from medical records of patients undergoing HD. Sampling was done using convenience sampling technique. A total of 130 files were reviewed from two sectors in Jordan. Data were collected from electronic patients' files.

**Results:** The result showed that erythropoietin stimulation agents' response as represented by hemoglobin level was negatively predicted from both, the Erythropoietin dose ( $B = -0.439$ ,  $p < 0.001$ ) and the iron level ( $B = -0.029$ ,  $p < 0.05$ ). While serum phosphorus ( $B = 0.002$ ,  $p = 0.958$ ), C-reactive protein ( $B = -0.001$ ,  $p = 0.574$ ), serum albumin ( $B = -0.027$ ,  $p = 0.405$ ) and vitamin B12 level ( $B = -3.07E-05$ ,  $p = 0.764$ ) didn't significantly predict the response to erythropoietin stimulation agents.

**Conclusion:** Frequently giving high doses of ESAs therapy may have a reverse effect on response to ESA therapy. Laboratory results should be frequently monitored. Additionally, iron level alone should not be relied on as an indicator for erythropoietin stimulation agents' responsiveness.

**Keywords:** Chronic Kidney Disease, Hemoglobin level, Hemodialysis patients, Erythropoietin Stimulating Agents, Jordan.